

FIRST RESULTS FROM THE ANGLO-AUSTRALIAN PLANET SEARCH – A BROWN DWARF CANDIDATE AND A 51 PEG-LIKE PLANET¹

C.G. TINNEY², R. PAUL BUTLER^{3,2}, GEOFFREY W. MARCY^{4,5}, HUGH R.A. JONES⁶, ALAN J. PENNY⁷, STEVEN S. VOGT⁸, KEVIN APPS⁹, GREGORY W. HENRY¹⁰

To appear in The Astrophysical Journal, Letters.

ABSTRACT

We report results from the Anglo-Australian Planet Search – a survey for planets around 200 solar-type stars in the southern hemisphere, which is being carried out on the 3.9m Anglo-Australian Telescope. Limiting Doppler precisions of 3 m s^{-1} have been demonstrated from the first 2.5 years of operation, making this the highest precision planet search in the southern hemisphere. From these data we report results for two new sub-stellar detections. The first is a “51 Peg”-like planet around the star HD 179949 with $M \sin i = 0.84 M_{\text{JUP}}$. Photometric study reveals this is not a transiting system. The second is a brown dwarf or very low-mass star companion to HD 164427 in an eccentric orbit with $M \sin i = 46 M_{\text{JUP}}$. Hipparcos data indicate this latter object is unlikely to have a mass greater than $0.18 M_{\odot}$.

Subject headings: planetary systems – stars: individual (HD164427, HD179949) – stars: low-mass, brown dwarfs

1. INTRODUCTION

Since the discovery of the first extra-solar planet by Doppler velocity techniques in 1994 by Mayor & Queloz (1995), planetary detections have been dominated by northern hemisphere search programmes – most prolifically by the precision velocity programmes at Lick (e.g. Butler et al. (1996)) and Keck (e.g. Vogt et al. (2000)), but also complemented by lower precision programmes at OHP (Baranne et al. 1996), McDonald Observatory (Cochran et al. 1997), AFOE (Noyes et al. 1997), and programmes at La Silla (Kürster et al. 2000; Queloz et al. 2000). Of these programmes, only the latter have access to the sky south of $\sim -20^\circ$, and these achieve precisions of $\sim 10 \text{ m s}^{-1}$. In 1998, therefore, the Anglo-Australian Planet Search was begun to complete the all-sky coverage of the brightest stars at precisions reaching 3 m s^{-1} . In this paper we present some first results from this programme. A companion paper (Butler et al. 2000, hereafter Paper II) presents results for a further two new planets, along with a detailed description of our observational programme.

2. THE ANGLO-AUSTRALIAN PLANET SEARCH

The Anglo-Australian Planet Search is being carried out on the 3.92m Anglo-Australian Telescope (AAT), using the University College of London Echelle Spectrograph (UCLES) and an I₂ absorption cell. UCLES is operated in its 31 lines mm^{-1} mode with an MIT/LL 2048×4096 $15 \mu\text{m}$

pixel CCD. This lumogen-coated CCD (denoted MITLL2) underwent a serious failure at the end of 1999, which required its read-out to be switched to the second of its working amplifiers in 2000 January (following which it was denoted MITLL2a). No change in the operation of the detector (apart from a flipped read-out format) has been detected by our programme.

Our target sample of ≈ 200 stars with $\delta < -20^\circ$, includes F,G and K V-IV stars with $V < 7.5$ and MV stars with $V < 11.5$. Where age/activity information is available from S or R_{hk} indices (see e.g., Henry et al. (1996)), target stars are required to have ages greater than 3 Gyr. Our first observing run was in 1998 January, and the last run for which observations are reported here was in 2000 November. The observing and data processing procedure follows that described in Butler et al. (1996), and is described in detail in Paper II. In particular, Paper II presents velocities for a number of stable stars illustrating that we reach a velocity precision floor of 3 m s^{-1} for the bright stars included in our programme. One of the stars presented here (HD 179949) has also had confirmatory observations acquired at three epochs in 2000 September 5-8, as part of the Keck planet search (Vogt et al. 2000).

3. CHARACTERISTICS OF HD 179949 & HD 164427

HD 179949 (HR 7291, HIP 94645, GJ749) is an F8V star, which shows moderate rotation with a measured $v \sin i = 6.3 \pm 0.9 \text{ km s}^{-1}$ (Groot et al. 1996). Its Hipparcos parallax puts it at $27.0 \pm 0.5 \text{ pc}$, with $M_V = 4.09 \pm 0.04$ (ESA

¹ Based on observations obtained at the Anglo-Australian Telescope, Siding Spring, Australia; and the W.M. Keck Observatory, which is operated jointly by the University of California and the California Institute of Technology.

² Anglo-Australian Observatory, PO Box 296, Epping. 1710. Australia. cgt@aaoepp.aao.gov.au

³ Carnegie Institution of Washington, Department of Terrestrial Magnetism, 5241 Branch Rd NW, Washington, DC 20015-1305

⁴ Department of Astronomy, University of California, Berkeley, CA, 94720

⁵ Department of Physics and Astronomy, San Francisco State University, San Francisco, CA 94132.

⁶ Astrophysics Research Institute, Liverpool John Moores University, Twelve Quays House, Egerton Wharf, Birkenhead CH41 1LD, UK

⁷ Rutherford Appleton Laboratory, Chilton, Didcot, Oxon OX11 0QX, U.K.

⁸ UCO/Lick Observatory, University of California at Santa Cruz, CA 95064.

⁹ Physics and Astronomy, University of Sussex, Falmer, BN1 9QJ, U.K.

¹⁰ Center of Excellence in Information Systems, Tennessee State University, Nashville, TN 37203-3401.

1997). There is no published evidence to indicate that HD 179949 is a binary. The photometry of Eggen (1998) indicates HD 179949 has roughly solar metallicity with $[\text{Fe}/\text{H}] = +0.02 \pm 0.1$. The *ubvy* calibration of Vogt et al. (2000) and published photometry would indicate a more metal-rich $[\text{Fe}/\text{H}] = +0.22 \pm 0.07$. The latter is supported by Twarog & Anthony-Twarog (1995, Fig. 8) which indicates HD 179949 is as metal-rich, if not more so, than the majority of disk main-sequence stars. HD 179949 has a moderate X-ray luminosity as determined by ROSAT’s PSPC – Hünsch et al. (1998) determine a $L_X = 41.0 \times 10^{27} \text{ erg s}^{-1}$, while in an independent analysis Piers et al. (1998) determine $L_X/L_{\text{bol}} = (9.7 \pm 3.3) \times 10^{-6}$. Both are some 10 times higher than the equivalent quantity in the quiet Sun. The spectra acquired at Keck have been used to derive a R'_{hk} index of -4.72 for this star, using the same procedure as Vogt et al. (2000). Figure 1 compares the Ca II H line with that seen in the Sun. The mass of HD 179949 is estimated to be $1.24 \pm 0.1 M_{\odot}$, based on interpolation between evolutionary tracks of Fuhrman et al. (1997, 1998).

HD 164427 (HIP 88531, Gl 700A) is an inactive G0V (Houck & Cowley 1975) star, with a R'_{hk} index of -4.95 (Henry et al. 1996). Its Hipparcos parallax puts it at a distance of $39.1 \pm 1.4 \text{ pc}$, and $M_V = 3.91 \pm 0.8$ (ESA 1997) making it somewhat over-luminous for its spectral type. Indeed Evans et al. (1964) classified it as a sub-giant, with luminosity class IV. The *ubvy* calibration of Vogt et al. (2000) suggests a metallicity of $[\text{Fe}/\text{H}] = +0.11 \pm 0.07$. HD 164427 was catalogued as a very wide binary (Gl 700AB) by Gliese (1969) (though both have since dropped out of the nearby star sample (Gliese & Jahreiss 1991) due to better parallaxes placing them outside the 25 pc limit). The binary identification is actually due to Luyten (1957) who catalogued this pair as LTT 7172 and LTT 7173 (respectively) with common proper motion and a $28''$ separation at a position angle of 336° . There is no subsequent astrometry of this pair in the literature. Examination of Digital Sky Survey scans of a UKST plate from 1992.6 shows no evidence for an object with the magnitude difference indicated by Luyten ($\Delta m = 7.2$) at this separation relative to HD 164427, leading us to conclude it may not be a common proper motion pair. Even if the system were a binary, it would be a very wide system (1090 au) – so wide as to be irrelevant for the purposes of high precision doppler velocities. The mass of HD 164427 is estimated to be $1.05 \pm 0.1 M_{\odot}$.

Both stars were seen to be photometrically stable over the life of the Hipparcos mission at a 95% confidence level of < 0.015 magnitudes (ESA 1997).

4. RADIAL VELOCITY OBSERVATIONS AND ORBITAL SOLUTIONS

Twenty-three observations of HD 179949 are listed in Table 1, where the column labelled “Uncert.” is the velocity uncertainty produced by the least-squares fitting process, which simultaneously determines the Doppler shift and the spectrograph point-spread function (PSF), given an iodine absorption spectrum, an “iodine free” template spectrum of the object, and an “iodine” spectrum of the object (Butler et al. 1996). This uncertainty includes the

effects of photon-counting uncertainties, residual errors in the spectrograph PSF model and variation in the underlying spectrum between the template and “iodine” epochs. Only observations where the uncertainty is less than twice the median uncertainty are listed. These data are shown in Figure 2 phased with a period of 3.093 days. The figure shows the best-fit Keplerian model for the data, with the resultant orbital parameters listed in Table 2. The best-fit period from Keplerian fitting is the same as the period found from the peak of the Scargle periodogram, with a false alarm probability less than 0.001. Both AAT (dots) and Keck (squares) data are shown, with the Keck data completely confirming the orbital fits derived from the AAT data alone. Like the previously observed giant planets in 3 to 5 day orbits, HD 179949 appears to be in a circular orbit. Table 2 includes the best-fit orbital solutions both for the case of a circular orbit and floating eccentricity. The resulting minimum companion mass is $0.84 M_{\text{JUP}}$, with an orbital semi-major axis of 0.045 au.

The residuals about the fit are at the level of “jitter” expected in an F8V star with HD 179949’s level of rotation (6.3 km s^{-1}) and activity ($R'_{hk} = -4.72$). “Jitter” here is used to refer to the scatter in the observed velocity about a mean value in systems observed over the long-term to have no Keplerian Doppler shifts. It is thought to be the combined effect of surface inhomogeneities, stellar activity and stellar rotation. Saar et al. (1998) have studied the correlations between “jitter”, and stellar rotation and activity in the Lick precision velocity programme. They indicate we would expect HD 179949 to show $\approx 10 \text{ m s}^{-1}$ “jitter” due to activity, spots, etc. We measure a scatter of 10 m s^{-1} about our Keplerian fit, which is consistent with this expectation.

The twenty-seven observations of HD 164427 are listed in Table 3, and they are shown in Figure 3 along with a Keplerian fit to the data with the orbital parameters listed in Table 2. The rms scatter about this fit is 7.7 m s^{-1} , slightly larger than the median uncertainty from the fitting process of 5.6 m s^{-1} . This is larger than the systematic precision limit of the Anglo-Australian Planet Search, as the earliest observations of this object were performed in very poor conditions. As soon as it was realised HD 164427 was a velocity variable with large amplitude, it was “prorated” within the programme to only short exposures (and so lower precision than the rest of the programme), though still more than sufficient to precisely determine the companion’s orbit. The resultant minimum companion mass is $46 M_{\text{JUP}}$, or $0.043 M_{\odot}$, and the orbital semi-major axis is 0.46 au.

5. DISCUSSION

5.1. HD 179949

Ten other “51 Peg”-like planets are currently known with periods under 10 days, and eccentricities less than 0.1¹¹. The minimum masses of these planets span the range $0.24\text{--}3.87 M_{\text{JUP}}$, with orbital semi-major axes in the range $0.038\text{--}0.059 \text{ au}$. In all respects, HD 179949 falls squarely into this class of objects. Even its above-solar metallicity is similar to that seen in the other “51 Peg”-like planets (e.g., Queloz et al. (2000), Gonzalez (2000)).

¹¹ In the absence of a formal classification scheme, we consider HD 83443, HD 168746, HD 46375, HD 75289, BD-10 3166, HD 187123, HD 209458, ν And b, τ Boo to be “51 Peg”-like, as well as 51 Peg itself.

5.2. Transit Search in HD 179949

As with all the other “51 Peg”-like planets, HD 179949 is an excellent candidate for a transit search. The Hipparcos magnitude, color index, and parallax of the star imply a radius of about $1.24 R_{\odot}$, which, combined with the orbital radius of 0.045 au, places the probability of observable transits at 12.8%. Successful transit observations (already carried out for HD 209458 by Charbonneau et al. (2000) and Henry et al. (2000b)) not only tightly constrain orbital inclination, but provide planetary radii as well.

To search for transits in HD 179949, we obtained photometric observations between 2000 September 21 and October 1 with the T8 0.80 m Automatic Photoelectric Telescope (APT) at Fairborn Observatory in southern Arizona¹². The instrumentation and techniques used for this transit search are described in Henry (1999) and Henry et al. (2000a). The observations were made differentially with respect to the comparison star HD 178075 ($V = 6.30$, $B - V = 0.01$, B9.5V). The precision of an individual observation is approximately 0.004 mag, somewhat worse than the typical precision of 0.001 mag obtainable with the APT due to observations being at air masses between two and three.

No transit events were detected at the times predicted by the orbital parameters of Table 2. For an assumed planetary radius of $1.4 R_{\text{JUP}}$ (the measured radius of the planetary companion of HD 209458 (Henry et al. 2000b; Jha et al. 2000)), the predicted transit depth is 0.014 mag. The mean of our 16 photometric observations of HD 179949 taken within the predicted transit window is 0.1058 ± 0.0010 mag; the mean of the 38 observations outside the transit window is 0.1044 ± 0.0007 mag. Thus, the two means agree within their respective errors and transits deeper than about 0.001 mag are ruled out. This non-detection of transits limits the orbital inclination i to less than 83° , and $\sin i$ to less than 0.992. Furthermore, the photometric observations place a limit of about 0.001 mag on any photometric variability of HD 179949 on the radial velocity period. This confirms that neither starspots nor stellar pulsations can be the cause of the radial velocity variations (see Henry et al. (2000a)) and so strongly supports the existence of the planet even in the absence of transits.

5.3. HD 164427

Recent work on the mass function of brown dwarfs in star clusters (Luhman et al. 2000; Lucas & Roche 2000; Zapatero Osorio et al. 2000) has failed to show that star formation processes cannot form objects below the $13 M_{\text{JUP}}$ deuterium burning limit. (Though the same results can be interpreted - less straightforwardly - as implying the existence of free-floating sub- $13 M_{\text{JUP}}$ objects formed by dynamical evolution ejecting objects formed by planetary processes.) Certainly they provide no evidence that the deuterium burning limit is a useful demarcation boundary between star- and planet-formation processes. Difficult as it may be, only an understanding of formation mechanisms (and possible subsequent dynamical evolution) can allow us to determine whether objects in the $5\text{--}20 M_{\text{JUP}}$ range are planets or brown dwarfs. Nonethe-

less, the minimum mass of the companion to HD 164427 clearly lies in a range unlikely to be due to planetary formation processes - it is a brown dwarf or a star, not a planet.

Hipparcos astrometry, however, allows us to place some limits on the orbital inclination and mass for HD 164427's companion. The astrometric solution for HD 164427 shows a good fit for the parallax and proper motion of a single star, with $1\text{-}\sigma$ residuals of ≈ 3 mas. The uncertainty on the derived parallax is 0.9 mas. Based on this we can assume an upper limit to the astrometric perturbation (a_0) of HD 164427 due to an unseen companion of ≈ 2 mas. Then from the relationship between the spectroscopic and astrometric orbital elements (e.g. Halbwachs et al. (2000), equation 1) we conclude that $\sin i > 0.24$, or equivalently that the companion mass is less than $190 M_{\text{JUP}}$ or $0.18 M_{\odot}$, making it either a brown dwarf, or a very low-mass star. With an orbital semi-major axis of 0.46 au at 39 pc, such a companion will have a maximum apparent separation of 11.8 mas. Table 4 compares the absolute K magnitudes of HD 164427, with predicted magnitudes of the companion in the range of allowed masses. Only at the very massive end of this range is the companion likely to be detectable by either ground-based adaptive optics or space-based imaging.

Nonetheless, such a challenging observation must be pursued. Of the small number of brown dwarf candidates identified by precision Doppler surveys to date (around 11), many have been shown to actually be stars (Halbwachs et al. 2000), making *bona fide* brown dwarf companions rare, and well worth unambiguously identifying.

6. CONCLUSIONS


We present first results from the Anglo-Australian Planet Search, which is now the highest precision Doppler planet search in the southern hemisphere. These include a new member of the “51 Peg”-like class of planets around the F8 dwarf HD 179949, and a new brown dwarf candidate companion to the G0 dwarf HD 164427. This programme has now been running on the AAT for over 2 years and has demonstrated velocity precisions reaching 3 m s^{-1} . We confidently expect many more discoveries in the years to come - particularly as it extends its sensitivity toward the Jupiter-like planet regime.


The Anglo-Australian Planet Search team would like to gratefully acknowledge the support of the Director of the AAO, Dr Brian Boyle, and the superb technical support which has been received throughout the programme from AAT staff - in particular E. Penny, R. Patterson, D. Stafford, F. Freeman, S. Lee, J. Pogson and G. Schafer. We further acknowledge support by; the partners of the Anglo-Australian Telescope Agreement (CGT, HRAJ, AJP); NASA grant NAG5-8299 & NSF grant AST95-20443 (GWM); NASA grant NAG5-4445 & NSF grant AST-9619418 (SSV); NSF grant AST-9988087 (RPB); NASA grants NCC5-96 & NCC5-228, NSF grant HRD-9706268 & the Richard Lounsbery Foundation (GWH); and Sun Microsystems.

¹² These observations are available at <http://schwab.tsuniv.edu/t8/hd179949/hd179949.html>.

REFERENCES

- Baranne, A., et al. 1996, A&AS, 119, 373
- Butler, R.P., Marcy, G.W., Williams, E., McCarthy, C. & Dosanji, P. 1996, PASP, 108, 500
- Butler, R.P., Tinney, C.G., Marcy, G.W., Jones, H.R.A., Penny, A.J., Apps, K. 2000, ApJ, submitted (Paper II)
- Chabrier, G., Baraffe, I., Allard, F. & Hauschildt, P. 2000, ApJ, in press astro-ph/0005557
- Charbonneau, D., Brown, T.M., Latham, D.W. & Mayor, M. 2000, ApJ, 529, L45
- Cochran, W.D., Hatzes, A.P., Butler, R.P. & Marcy, G.W. 1997, ApJ, 483, 457
- ESA, 1997, The Hipparcos and Tycho Catalogues, ESA SP-1200
- Eggen, O.J. 1998, AJ, 115, 2397
- Evans D.S., Laing J.D., Menzies A. & Stoy R.H., 1964, R.Obs.Bull, 85, 207
- Fuhrmann, K., Pfeiffer, M.J. & Bernkopf, J. 1997 A&A, 326, 1081
- Fuhrmann, K., Pfeiffer, M.J. & Bernkopf, J. 1997 A&A, 336, 942
- Gliese, W. 1969, Catalogue of Nearby Stars, Edition 1969, Verlag G.Braun: Karlsruhe.
- Gliese, W. & Jahreiss, H. 1991, Preliminary Version of the Third Catalogue of Nearby Stars, Astronomischen Rechen-Institut: Heidelberg.
- Gonzalez, G. 2000, ASP Conf. Ser. (in press), Disks, Planetesimals, and Planets, ed. F. Garzon, C. Eiroo, D. de Winter & T.J. Mahoney, (San Francisco: ASP)
- Groot, P.J., Pters, A.J.M. & van Paradijs, J. 1996, A&AS, 118, 545
- Halbwachs, J.L., Arenou, F., Mayor, M., Udry, S. & Queloz, D. 2000, A&A, 355, 581
- Henry, T.J. & McCarthy, Jr, D.W. 1993, AJ, 106, 773
- Henry, T.J., Soderblom, D.R., Donahue, R.A. & Baliunas, S.L. 1996, AJ, 111, 439.
- Henry, G.W. 1999, PASP, 111, 845
- Henry, G.W., Baliunas, S.L., Donahue, R.A., Fekel, F.C. & Soon, W. 2000, ApJ, 531, 415
- Henry, G.W., Marcy, G.W., Butler, R.P. & Vogt, S.S. 2000, ApJ, 529, L41
- Houck, N. and Cowley, A.P. 1975, Michigan Catalogue of Two Dimensional Spectral Types for the HD stars, Volume 1, Michigan Spectral Survey: Ann Arbor
- Houck, N. and Smith-Moore, M. 1988, Michigan Catalogue of Two Dimensional Spectral Types for the HD stars, Volume 4, Michigan Spectral Survey: Ann Arbor
- Hünsch, M., Schmitt, J.H.M.M. & Voges, W. 1998, A&AS, 132, 155
- Jha, S., Charbonneau, D., Garnavich, P.M., Sullivan, D.J., Sullivan, T., Brown, T.M. & Tonry, J.L. 2000, ApJ, 540, L45
- Kürster, M., Endl, M., Els, S., Hatzes, A. P., Cochran, W. D., Döbereiner, S. & Dennerl, K. 2000, A&A, 353, L33
- Lucas, P.W. & Roche, P.F. 2000, MNRAS, 314, 858.
- Luhman, K.L., Rieke, G.H., Young, E.T., Cotera, A.S., Chen, H., Rieke, M.J., Schneider, G. & Thompson, R.I. 2000, ApJ, 540, 1016
- Marcy, G.W. & Butler, R.P. 1998, ARA&A, 36, 57
- Mayor, M. & Queloz, D. 1995 Nature, 378, 355
- Noyes, R., Jha, S., Korzennik, S., Krockenberger, M., Nisenson, P., Brown, T., Kennelly, E. & Horner, S., 1997, ASP Conf.Ser., 119, 119
- Pters, A.J.M., van Paradijs, J. & Schmitt, J.H.M.M. 1998, A&AS, 128, 29
- Queloz, D. et al., 2000, A&A, 354, 99
- Saar, S.H., Butler, R.P. & Marcy, G.W. 1998, ApJ, 403, L153
- Twarog, B.A. & Anthony-Twarog, B.J. 1995, AJ, 109, 2828
- Vogt, S.S., Marcy, G.W., Butler, R.P. & Apps, K. 2000, ApJ, 536, 902
- Zapatero Osorio, M.R., Béjar, V.J.S, Martín, E.L., Rebolo, R., Barrado y Navascués, D., Bailer-Jones, C.A. & Mundt, R., 2000, Science, 290, in press








TABLE 1
VELOCITIES FOR HD 179949

JD (-2440000)	RV (m s^{-1})	Uncert (m s^{-1})
11120.910	118.1	5.8
11383.024	-43.3	8.9
11410.943	-47.7	5.3
11413.052	-15.0	4.7
11413.924	-42.1	6.1
11472.917	-3.6	5.7
11683.133	-40.6	8.0
11684.180	113.7	6.1
11706.151	91.1	9.8
11718.161	127.1	6.8
11742.960	115.9	5.9
11743.992	-42.9	5.9
11745.101	0.0	5.8
11766.970	40.8	6.2
11768.000	109.6	8.5
11770.107	44.6	6.1
11792.751 ^a	122.2	2.6
11793.798 ^a	-80.7	2.5
11795.803 ^a	115.6	2.5
11827.956	-74.3	6.1
11828.911	61.5	6.7
11829.915	117.1	8.2
11855.942	-74.4	10.3

^aObservations from Keck. All other observations are from the AAT.

TABLE 2
ORBITAL PARAMETERS

Parameter	HD 179949 ($e=0.0$)	HD 179949 (e fitted)	HD 164427
Orbital period P (d)	3.093 ± 0.001	3.093 ± 0.001	108.55 ± 0.04
Velocity amp. K (m s^{-1})	101.3 ± 3.0	102.2 ± 3.0	2229 ± 77.0
Eccentricity e	0.0 (fixed)	0.05 ± 0.03	0.55 ± 0.02
ω ($^\circ$)	0.0	226 ± 25	356.9 ± 0.5
$a_1 \sin i$ (km)	4307.3 ± 15.0	4338.7 ± 15.0	$(2.776 \pm 0.049) \times 10^6$
Periastron Time (JD-244000)	11718.19 ± 0.1	11723.24 ± 0.1	11724.6 ± 0.2
$M \sin i$ (M_{JUP})	0.84 ± 0.05	0.84 ± 0.05	46.4 ± 3.4
a (AU)	0.045 ± 0.004	0.045 ± 0.004	0.46 ± 0.05
RMS about fit (m s^{-1})	10.8	10.3	7.7

TABLE 3
VELOCITIES FOR HD 164427

JD (-2440000)	RV (m s^{-1})	Uncert (m s^{-1})
10917.287	-1071.2	7.5
11002.091	-897.7	10.2
11118.891	-1026.5	9.8
11119.905	-1025.7	8.3
11236.281	-1065.7	6.4
11274.300	-3.9	5.3
11385.868	347.2	6.8
11386.879	513.8	4.9
11410.911	703.9	4.2
11413.034	378.0	3.6
11413.904	286.7	5.6
11472.893	-914.8	4.6
11630.300	391.3	4.9
11683.096	-1014.7	5.6
11684.147	-1006.3	6.4
11706.112	-217.9	4.9
11717.923	1718.7	5.3
11718.123	1775.8	4.9
11742.913	-44.6	5.5
11743.933	-126.6	7.0
11745.067	-218.7	4.6
11766.942	-968.3	5.2
11767.992	-985.4	8.4
11827.914	2169.0	5.5
11828.890	2480.8	6.3
11829.894	2791.5	7.0
11855.921	-371.0	8.0

TABLE 4
PREDICTED K MAGNITUDES FOR HD 164427 COMPANION

Companion Mass	Age	M_K	ΔM_K
HD 164427 $M_K=5.4$			
$0.18 M_\odot^a$	...	7.7	2.3
$0.10 M_\odot^a$	...	9.2	3.8
$75 M_{\text{jup}}^b$	1 Gyr	11.2	5.9
	5 Gyr	11.7	6.3
$42 M_{\text{jup}}^b$	1 Gyr	13.7	8.3
	5 Gyr	>16	>10.6

^aEmpirical M_K values due to Henry & McCarthy (1993).

^bModel M_K values due to Chabrier et al. (2000).